

Pediatric Neurosurgery in Pakistan: Current Status and a Call-To-Action for Inclusion in The National Surgical Plan

Samaa Akhtar MD, MHSc¹, Javaid Iqbal MBA, MA, MS²

1. Pakistan Children's Heart Foundation
2. Brandeis University

Abstract:

The pediatric population in Pakistan is vulnerable to premature morbidity and mortality due to ineffective neurosurgical health system performance. Approximately 7 per 1000 children in Pakistan have epilepsy, cerebral palsy, and central nervous system infections. The ramifications are dire when this incidence rate is paired with a scarcity of skilled pediatric neurosurgeons, resources, and infrastructure. Traumatic Brain Injuries (TBI) are one of the leading causes of death in children under three years of age. 76% of which are avertible with timely surgical access. The National Surgical, Obstetric, and Anesthesia Plan (NSOAP) provides a framework to facilitate access to surgical treatment and improve surgical system resilience. This paper presents a commentary on the status of pediatric neurosurgery in Pakistan and a seminal framework for policy reform to advocate its inclusion in the NSOAP.

KEYWORDS: *Pediatric; Neurosurgery; NSOAP; Pakistan; GlobalSurgery*

Corresponding author:

Samaa Akhtar MD, MHSc

Pakistan Children's Heart Foundation

Email: Samaa.akhtar@pchf.org.pk

Introduction

The global burden of neurological disorders is approximately 6.5%¹. In low-income countries (LICs), the prevalence rate ranges between 5 to 8.4% compared to a range of 10-11% in high-income countries (HICs)¹. LICs report higher mortality and morbidity ratios than HICs. The principal reason is the unavailability and inaccessibility of timely neurosurgical treatment by the affected population. Safe and affordable surgery is not accessible to approximately 1.7 billion children worldwide. Considering that 40% of the population in LICs is under the age of 15, the ramifications are catastrophic regarding Survival and Disability Adjusted Life Years (DALY) and death². Currently, there is insufficient global and country-wide data on the age-standardized incidence, prevalence, and mortality rates specific to neurological disorders in LICs in particular. In a global context, the pediatric neurosurgical burden worldwide is alarming. Approximately 400,000 new cases of hydrocephalus are recorded each year³. The incidence of hydrocephalus is reported as 123 per 100,000 births in low- and middle-income countries (LMICs) alone⁴. The incidence of epilepsy and traumatic brain injury (TBI) in children is estimated to be 1.4 million and 3 million, respectively⁴. Pakistan has a population of around 207 million. 35% are below the age of 15 years⁵.

It is, therefore, alarming that there are currently no national surveys on the epidemiological data of pediatric neurology and neurosurgery in Pakistan. Most surveys, research, and reports can be sourced from hospital-based studies. According to one survey, 4-5% of Pakistan's non-communicable diseases (NCDs) are neurological disorders⁶. Other developing

countries, such as India, report a mean prevalence rate of neurological conditions as 2,394 per 100,000 population⁶. There is no data on the prevalence rate particular to the pediatric population, however. Disease-specific surveys, however, report that 7 per 1000 children in Pakistan have epilepsy, cerebral palsy, and central nervous system infections⁶. Health indicators in Pakistan also remain high. The under-five mortality rate is 94 per 1000 births. The Infant Mortality Rate (IMR) is 78 per 1000 live births. Primary causes of mortality include diarrheal diseases and acute respiratory infections (ARI)⁷. There are no national surveys on the ratio of IMR attributed to pediatric neurosurgical conditions. However, Waqas et al.⁸ reports head injuries in infants as a predominant cause of in-hospital mortality, at 7.3%. 76% of these head injuries are preventable through improved parental safety measures⁸.

Unfortunately, the vast need for understanding and improving the pediatric neurosurgery infrastructure in Pakistan has remained relatively unexplored and understudied. For this reason, the 'Global Surgery 2030' report by The Lancet Commission on Global Surgery (LCoGS) was a much-needed step forward⁹. The development of the National Surgical, Obstetrics, and Anesthesia Plans (NSOAPs) has brought surgery to the table for national health policy reform. Pakistan and Nigeria are among the forerunners in integrating the NSOAP process into their National Surgical Plans and incorporating pediatric surgery into the agenda¹⁰. There are several challenges ahead, however. Pediatric neurosurgical diseases should be incorporated to reach the NSOAP target of 80% essential anesthesia and surgical services coverage⁹.

Pediatric neurosurgery is grossly underrepresented in Pakistan. Access to timely and quality treatment for patients and accurate data for crucial health stakeholders and policymakers is severely compromised. Acute surgical illnesses alone result in 187 deaths per 100,000 population in Pakistan⁹. Death and disability in pediatric neurosurgery only add to the high IMR. It is essential to conduct neurological epidemiological studies in Pakistan. To identify and address the critical deficits in knowledge, policy, and action. All factors considered, there is a dire need for Pakistan's pediatric neurosurgical sector to be included in the National Surgical Plan. Inclusion in the National Surgical Plan will also help financing options for pediatric neurosurgery under the recently introduced Universal Health Coverage Program, known as The Sehat Sahulat Program¹¹. Currently, only general neurosurgeries are covered under this program¹¹. Amending policies to include pediatric neurosurgery cover will be a substantial step towards sustainability.

Pakistan's Healthcare System

Pakistan has a two-tier healthcare system. There are approximately 920 hospitals in the public sector and around 800 in the private sector. Bringing the estimated total number of beds to 100,000 only⁷. Being the sixth most populated country in the world, Pakistan is one of 57 countries with a critical deficit in health workforce and resource deficiency⁷. Only 2.8% of Pakistan's gross domestic product (GDP) goes to health expenditure (12). The per capita health expenditure is US\$37, and the government covers only 37%. Approximately 78.8% of healthcare financing is dominated by out-of-pocket payments¹². The remaining is covered by government taxation and

revenues, private insurance, and global funding. However, only 2% of foreign aid is allocated for health⁷. Medical insurance cover is rudimentary, which makes financing surgeries a considerable challenge in particular. Approximately 50% of Pakistan's population resides below the poverty line, that is, US\$2 per day per capita income¹², which is catastrophic for the financial risk protection of the population. Where covering basic medical expenses proves difficult, affording technical neurosurgical procedures seems impossible for patients. The socio-economic and logistic disparity also present challenges for the rural community, in particular, to access health. Difficulties in referrals to tertiary hospitals due to the unavailability of transport facilities are shared. In an attempt to curb this, there are around 5336 basic health units, 595 rural health centers, and 1138 mother and child centers nationwide⁷. Owing to poor management, however, the provision of adequate healthcare in these centers is ineffective. The most significant challenges Pakistan's health system faces are poverty, an increase in the burden of non-communicable diseases where communicable diseases are already prevalent, and a lack of quality health services. Even though there has been improvement, lessons should be taken from the failures of achieving the Millennium Development Goals of 2015 in order to achieve the targets of the surgical vision of 2025⁷.

The Pediatric Neurosurgery Infrastructure in Pakistan

Neurosurgery was established as a distinct surgical discipline only six decades ago in Pakistan. The first center was established by Omar Vali Jooma, in Karachi, at the Jinnah

postgraduate medical center in 1951¹³. Expansion followed shortly after. The Combined Military Hospital, Lahore, was established as the second center by Brigadier G.D. Qazi in 1955. In 1963, Bashir Ahmad established three more centers across Pakistan's second largest province, Punjab. These included Nishtar Medical College and Hospital, Multan, King Edward Medical College, and Lahore General Hospital¹³.

There are around 42 neurosurgery centers across Pakistan, 150 practicing neurosurgeons, and approximately 1500 neurosurgery beds¹³. Resource allocation toward pediatric neurosurgery is compromised due to the lack of pediatric neurology-specific facilities. There are only two specialized neurology and spine centers in the country. The largest and most well-equipped neurology healthcare center is the Punjab Institute of Neurosciences¹⁴. It is a 500-bed hospital with an output of approximately 7000 neurosurgeries per annum. However, there is no data on the percentage allocation of beds and services to pediatric patients. Nor the pediatric patient turnover rate. Neuro-diagnostics is equipped with only 3 CT scans, 2 MRIs, and multiplanar angiography¹⁴.

There is no hospital in the country exclusive to pediatric neurology or neurosurgery. Pediatric neurosurgery is a part of multidisciplinary tertiary care hospitals. Most of these are in the public sector. Renowned hospitals include the National institute of child health (Karachi), Children hospital (Karachi), the Institute of Child Health and Children's Hospital (Lahore), Children Hospital Complex and Institute of Child Health (Multan), National Institute of Child Health (Karachi) and Children's Hospital (Quetta). Pediatric neurosurgery is also catered to in general hospitals and

private neurology clinics. However, on average, there are only 40 beds allocated for pediatric neurosurgery patients in such settings¹⁵.

Pakistan's division of neurosurgeons to the population is 1/1,130,000, whereas the global recommended ratio is 1/230,000¹³. While this places Pakistan ahead of Africa at 1/ 1,352,000, a stark difference can be seen with Europe at 1/121,000 and North America at 1/81,000¹³. This disproportion causes neurosurgery residents to manage the patient load of pediatric neurosurgery without supervision. Subsequently affects patient safety, the quality of surgeries performed, and health outcomes. Some of the more innovative neurosurgical options are available to children in Pakistan. These include deep brain stimulation (DBS), minimally invasive spine surgery, and endoscopic tumor and eyebrow keyhole brain surgery. These procedures are, however, performed by general neurosurgeons. Pediatric neurosurgeons primarily cater to standard procedures- such as the third ventriculostomy for hydrocephalus, craniosynostosis, and myelomeningocele correction¹⁶.

At present, there is no national epidemiological data on pediatric neurology and neurosurgery in Pakistan. A handful of studies report institution-based data. In Wasay et al. ¹⁷, a survey was conducted for 46021 patients on the prevalence of neurological disorders in a tertiary care hospital. The most common condition was benign epilepsies syndrome (33%), and the least common was developmental disorders (4.8%)¹⁷. Hussain et al. corroborates this data¹⁸.




Figure 1. Predominant Pediatric Neurosurgical Conditions in Pakistan According to a Study-Based Survey (18).

Epilepsy

The prevalence of epilepsy in children in Pakistan is around 15.5 to 23 per 1000 children¹⁹. It is exceptionally high in children in the rural population. Treatment disparity is such that 27.5% of patients in urban areas are treated, whereas only 1.9% of patients in rural areas. The etiology is idiopathic in 21 to 76% of cases¹⁹. According to Jawaid et al.²⁰, the most common causes are neonatal encephalopathy with an incidence rate of 28.3%, CNS infections at 19.3%, and febrile seizures at 12.7%. Secondary seizures are more common compared to primary seizures. Around 75.4% of children suffer from generalized onset tonic-clonic seizures, 5.7% from mixed seizures, and 5.2% from focal or bilateral tonic-clonic seizures²⁰.

Treatment options are limited due to the high cost of surgery. However, the cost of neurosurgery epilepsy in Pakistan is comparatively less than compared to countries like Chile and India²¹. A pediatric anterior temporal lobectomy costs around US\$1644 in the public sector and US\$3044 in the private sector. This procedure costs around US\$5,020 in Chile and US\$5000 in neighboring India²¹. Neurosurgical options are, however, limited. Vagus Nerve Stimulator (VNS) surgery for refractory epilepsy is almost non-existent in Pakistan. There has been only one reported case of a 7-year-old patient²². The procedure was successful and encouraged the need for neurosurgeries to help improve the quality of life for children with epilepsy in Pakistan. Considering the extrapolated number of epileptic children waiting for surgery is

around 34,000-45,000, the limitation for treatment is dire²³.

Cerebral palsy

Cerebral palsy (CP) is one of the most common motor disabilities in children in Pakistan²⁴. This burden is due to the high incidence of birth trauma and asphyxia, low birth weight (LBW) in neonates, and maternal and intracranial infections like meningoencephalitis and kernicterus in the region. CP is more prevalent in females and affects children with a mean age of 5.6 ± 2.25 years²⁴. Spastic quadriplegic CP is the most prevalent type due to the high incidence of birth asphyxia and trauma in the region. This is followed by atonic, ataxic/mixed CP, and finally athetoid CP. In terms of etiology, birth asphyxia in premature infants results in a high incidence of spastic diplegic CP, LBW, kernicterus, and intracranial hemorrhage contribute to a high incidence of spastic hemiplegic CP²⁴. Even though procedures like intrathecal baclofen pump surgery are available, no evidence suggests that it has been applied to the pediatric population in Pakistan²⁵. Other surgical options, such as selective dorsal rhizotomy, have only recently become available in Pakistan²⁵. That too in a private setting, with limited accessibility due to long waiting times and limited surgeons.

Hydrocephalus

In a survey by Asif et al.²⁶, 70% of hydrocephalus patients in a sample of 200 patients were congenital, while 30% were acquired. Spinal dysraphism accounted for 40% of congenital hydrocephalus. The most common causes of acquired hydrocephalus were post-meningitis (20%), brain tumors (5%), and post-traumatic (3%). The

neoplastic group consisted of 10 patients. 8 of these had posterior fossa tumors, and two had craniopharyngiomas. 50% of patients with congenital hydrocephalus had associated congenital anomalies.

Common post-operative complications included shunt obstruction (24%), infection (20%) intracerebral hemorrhage (0.5%) and subdural hematoma (0.5%)²⁷. Such postoperative complications are more common in low-socioeconomic settings like Pakistan. Fundamental reasons include late follow-up presentation of patients, late intervention, and lack of aseptic surgical conditions. Many patients do not seek timely medical attention because of social taboos²⁸. At the same time, others are intimidated by the cost of treatment. According to this study, the rate of shunt blockage in congenital and post-meningitis patients was also high. Common blockage causes included shunt infection (14%) and shunt failure (10%). 50% of blockages occurred within the first year of surgery. The most common type of shunts used in Pakistan are ventriculoperitoneal (VP), medium pressure of 50-100 mm shunts²⁷.

The use of an endoscopic third ventriculostomy (ETV) shunt for obstructive hydrocephalus is considered a routine procedure with a 30-90% success rate²⁹. It is, however, not as commonly used in Pakistan. Even though the comparative cost with VP shunt is almost similar at US\$ 2,177³⁰. One reason could be the lack of availability of the ETV shunt or the lack of skilled pediatric neurosurgeons available to perform the procedure. ETV shunts require more technical expertise in maneuvering the endoscope and the endoscopic anatomy³⁰. VP shunts also prove more expensive in the long run due to multiple surgeries needed to correct postoperative complications²⁹.

Brain tumors

Pediatric brain tumors are one of the most common causes of oncology-related mortality and morbidity globally and in Pakistan. They present added challenges for the pediatric population in Pakistan, however. There is no central pediatric tumor registry in Pakistan. The paucity of data and weak neurosurgery infrastructure leads to delayed diagnosis and treatment. These factors affect the prognosis of patients, even in uncomplicated cases. While nationwide data is not available, Riaz et al.³¹ is a single-center study that reports the clinical data of primary CNS tumors in children from 0-16 years between 2004 to 2014. From a sample of one hundred seventy-five children, glial tumors were the most common at 105 (60%). Embryonal tumors were at 40 (22.9%), craniopharyngiomas at 25 (14.3%), and germ cell tumors 1 (0.6%)³¹. The most common glial tumor was astrocytoma (25.7%), and the medulloblastoma (15.4%) was the most common embryonal tumor. Mortality was at 24%, while 11.4% of patients went into remission³¹.

Pediatric TBI

According to Khan et al.³², the incidence rate of TBI in Pakistan is 50/100,000 per annum. 73% of these injuries occur in male children under the age of 12. In another survey by Wani et al.³³, the incidence of 'falls' was 63% in pre-school children. Inferring falls in children is the most common cause of pediatric TBI. 25% of TBIs in children were caused by road traffic accidents (RTAs), and 11.6% were caused by other forms of injury, such as sports or collisions. Only about 30% of children had moderate to severe head injuries, and a Glasgow Coma Scale (GCS) score between 9-12³³. Most of the children

in these categories had poor surgical outcomes. Common injuries ranged from 10% of children having depressed fractures and contusions, 7.5% of children having extradural hematomas, 5% having subdural hematomas with brain contusions, and 6.66% of children having brain edema³³. 7.5% of patients underwent craniotomies and hematoma evacuation, and 10% underwent dural repairs and elevation of depressed fractures. Only 5% of children underwent decompression craniotomies³³. It is noteworthy to mention that according to the survey, 30% of deaths were avertible. Delays in surgery caused by long waiting times increase mortality by enhancing secondary injuries to the brain, such as hypotension and hypoxia³³.

Challenges

Pediatric neurosurgery is a particular niche and a technical specialty. In Pakistan, it is a part of multidisciplinary tertiary care hospitals, which results in gross differences between departments. In terms of skilled professionals, resource allocation, budgetary preference, and infrastructure. General neurosurgeons and pediatric surgeons perform surgeries such as VPS and spinal dysraphism instead of pediatric neurosurgeons. As a result, pediatric neurosurgeons do not cover even high-risk, specialized surgeries such as detethering an asymptomatic cord and the excision of intramedullary spinal cord tumors. Improved coordination between pediatric neurosurgeons, neurologists, and pediatricians can help navigate things more efficiently but cannot cover the overall shortage³⁴. Covering the skilled professionals' deficit is also challenging because of inadequate training programs for pediatric neurosurgery trainees. The existing

panel of pediatric neurosurgeons and trainees struggle to manage the patient volume as it is. It is inadvertently compromising the quality of care for patients. Additionally, there are no regularized morbidity meetings or undertaking of cutting-edge research projects and presentations to help improve the existing standard of care¹³.

Infrastructure issues are predominantly related to the basic infrastructure of hospitals in LICs. According to a World Health Organization (WHO) survey of 800 facilities in LICs, 31% of hospitals did not have continuous electricity, 22% did not have running water, and 24% did not have oxygen cylinders³⁵. Essential services such as accidents and emergency care, postoperative care, pathology, and blood transfusion services are also compromised³⁵. Corruption is also prevalent in the healthcare sector. It manifests itself in the form of bribes for preferential treatment and resource allocation within specialties.

According to Haque et al.², most pediatric departments in Pakistan are equipped with essential equipment, incubators, and phototherapy equipment. However, most pediatric Intensive Care Units and operation theatres do not have vital equipment such as ventilators, anesthetic equipment, and patient monitors². There is a lack of availability of designated pediatric intensive care units (PICU). For around 500,000 children, there is only 1 PICU bed per ICU. In comparison, the allocation ratio is one bed/10,000 children in the USA and 1bed/1000 in the UK². The availability of pediatric neurosurgical procedures is, however, also limited. Awake craniotomy or minimally invasive surgery is not even an option for children in Pakistan. Sub-disciplines such as functional neurosurgery, epilepsy surgery, and stereotactic

radiosurgery are almost non-existent. There is a shortage of neuronavigational facilities and endoscopy. To overcome this deficit, departments such as radiology, ENT, ophthalmology, and anesthesia support surgery coordination as much as possible¹³. Another example includes the limited availability of autologous bone grafts for pediatric cranioplasties. It is a better choice due to better osseointegration in the skull and easy storage. In Pakistan, however, neurosurgery is adapted to manage the burden of low resources. During craniectomy, the ovariectomized bone flap is inserted into the abdominal subcutaneous fat instead³⁶.

Neurosurgical emergency care and admissions are delayed extensively due to the lack of protocols for pre-hospital services. The absence of implementation of postoperative management in ICUs reduces patient turnover. Lack of transport, such as ambulances, is a debilitating aspect for TBI patients. If the TBI is complicated or severe, emergency care often refuses to take the patient because of a lack of skilled personnel for surgical treatment. The patient usually dies waiting.

In terms of equipment, there is only one tesla MRI machine in the public sector in Pakistan³⁷. Including DTI imaging, which has proven essential for tumor resection. There is also only one reported Cavitron ultrasonic surgical aspirator (CUSA)³⁷. There are added challenges for the rural population, such as diagnosing conditions. Patients have to travel long distances just for a simple X-ray. In Pakistan, it is also common practice for the late presentation of pediatric neurosurgical patients to neurosurgeons. Cultural stigmas, fear of the cost of treatment and the treatment itself, and lack of knowledge of the disease and treatment options are common reasons.

Implications for Advancing Pediatric Neurosurgery Towards the Sustainable Development Goals

A forward trajectory should involve minimizing losses and making the current pediatric neurosurgical infrastructure more efficient. For example, large-scale surveys can help identify key problem areas. Data from these surveys can allow targeted investment for the most impact generated with limited economic buy-in. International stakeholder engagement should also include incorporating the niche of the targeted population, that is, children. Such as The Global Initiative for Children's Surgery and Global PaedSurg³⁸. A similar approach should be applied to pediatric neurosurgical capacity building. The frequency of training programs and international medical missions should be increased. For example, the Foundation for International Education is currently running projects such as the Tanzania Neurosurgery Project, the Duke East Africa Neurosurgery Program, and the Cambodia Neurosurgery Support Project³⁹. Programs such as The World Federation of Neurosurgical Societies (WFNS) provide training for early career neurosurgeons in LICs³⁹.

Similar programs can be put together in Pakistan. For retention of the workforce,

learning should be competency-based with appropriate incentives. Apart from this, the standardization of protocols can be a low-cost driver to help reduce system losses and increase accountability. According to Fezeu et al.⁴⁰, there is an array of mapping techniques for epilepsy. There is also no consensus on a fixed protocol of anesthesia in awake craniotomies⁴⁰. Non-standardization can affect the best neurosurgical approach to addressing the condition. Service delivery can be improved through electronic statistical records. Particularly in computerized neurology charts, which are lacking in Pakistan. These tools can also improve multidisciplinary communication within tertiary care hospitals. Coordination between neurosurgeons and pediatricians, for example. Financing solutions should include minimizing recurrent losses. For example, in Uganda, the cost per DALY averted for neurosurgical intervention ranged from \$59 to \$126 in a survey⁴¹. The economic benefit incurred for treating hydrocephalus alone ranged from \$930 million to \$1.6 billion⁴¹. Pakistan can implement such strategies by minimizing shunt infections and blockages, providing timely surgeries, and increasing the use of antibiotic-impregnated hydrocephalus shunts.




Figure 2. A Seminal Framework Incorporating Pediatric Neurosurgery with the PSOAP Framework

Conclusion

The inherent need to include this vital subspecialty in the NSOAP is reiterated in examining the current status of pediatric neurosurgery in Pakistan. Engaging local leaders and stakeholders can provide a comprehensive and collaborative approach. The first step in adopting a decentralized NSOAP framework to a provincial model (PSOAP) has already laid the groundwork for a positive trajectory. This model can promote public-private partnerships within provinces. Provincial competition for funding is also reduced as stakeholder engagement is based on shared interests in investment. Which can subsequently help drive policies and legislation to bring

sustainability to the pediatric neurosurgical infrastructure in the region.

Standardized protocols for quality management can help streamline processes and policy frameworks. Since pediatric epilepsy and cerebral palsy are the predominant neurosurgical disorders, addressing the following patient population can help alleviate the patient volume. Surgical outcomes should also be evaluated and monitored to set a precedent for future guidelines. Health financing models should be adapted to minimize losses incurred. Bridging weaknesses in the current infrastructure and utilizing a strategic reorganization protocol can help include pediatric neurosurgery in the national surgical program in Pakistan.

References

1. Hussain, G., Shahzad, A., Anwar, H., Mahmood, Baig, S., Shabbir, A., & De Aguilar, J. L. G. (2017). Neurological disorder burden in Faisalabad, Punjab-Pakistan: data from the major tertiary care centers of the city. *Pakistan Journal of Neurological Sciences (PJNS)*, 12(3), 3-10.
2. Haque, A., Ladak, L. A., Hamid, M. H., Mirza, S., Siddiqui, N. R., & Bhutta, Z. A. (2014). A national survey of pediatric intensive care units in Pakistan. *Journal of Critical Care Medicine*, 2014.
3. Dewan, M. C., Rattan, A., Mekary, R., Glantz, L. J., Yunusa, I., Baticulon, R. E., ... & Warf, B. C. (2018). Global hydrocephalus epidemiology and incidence: systematic review and meta-analysis. *Journal of neurosurgery*, 130(4), 1065-1079.
4. Dewan, M. C., Baticulon, R. E., Rattani, A., Johnston, J. M., Warf, B. C., & Harkness, W. (2018). Pediatric neurosurgical workforce, access to care, equipment, and training needs worldwide. *Neurosurgical focus*, 45(4), E13.
5. Fatima I, Shoman H, Peters AW, Samad L, Nishtar S. Pakistan's national surgical, obstetric, and anesthesia plan: an adapted model for a devolved federal-provincial health system. *Canadian Journal of Anesthesia/Journal Canadien d'anesthésie*. 2020 Sep;67(9):1212-6.
6. The Aga Khan University. (2020). A dire need for trained pediatric neurologists in Pakistan. Retrieved 2021, from www.aku.edu/news/Pages/News_Details.aspx?nid=NEWS-002098.
7. Kumar, S., & Bano, S. (2017). Comparison and analysis of health care delivery systems: Pakistan versus Bangladesh. *J Hosp Med Manage*, 3(1), 21-22.
8. Waqas M, Jooma R. Unintentional head injury in children under three years of age: an insight into safety practices in a developing country. *Pediatric neurosurgery*. 2017;52(5):306-12.
9. Fatima, I., Shoman, H., Peters, A. W., Samad, L., & Nishtar, S. (2020). Pakistan's national surgical, obstetric, and anesthesia plan: an adapted model for a devolved federal-provincial health
10. Abbas A, Samad L. Children at the heart of global surgery: children's surgery in low-and middle-income countries.
11. Health Program PM. Sehat Sahulat Program [Internet]. About the Program. 2021 [cited 2021]. Available from: <https://www.pmhealthprogram.gov.pk/about-us/>
12. Organization WH. Pakistan [Internet]. Health Financing. World Health Organization; 2021 [cited 2021Sep]. Available from: <http://www.emro.who.int/pak/programmes/health-financing.html>
13. Shamim, M. S., Tahir, M. Z., Godil, S. S., Kumar, R., & Siddiqui, A. A. (2011). A critical analysis of the current state of neurosurgery training in Pakistan. *Surgical neurology international*, 2.
14. Institute, P. M. (2021). Department of Neurosurgery. Department of Neurosurgery | Postgraduate Medical Institute, Lahore. Retrieved September 2021, from http://pgmipunjab.edu.pk/dept_neurosurgery#3.
15. 14. Pakistan, I. R. of. (2018). DATA COLLECTION SURVEY ON HEALTH FACILITIES AND EQUIPMENT IN THE ISLAMIC REPUBLIC OF PAKISTAN FINAL REPORT (pp. 1–116). Islamabad, Punjab: Japan International Cooperation Agency (JICA).
16. Raza A. First-ever surgeries to treat epilepsy done in Pakistan [Internet]. Home Today's Paper National First-ever surgeries to treat epilepsy done in Pakistan. *The News*; 2021 [cited 2021Sep]. Available from: <https://www.thenews.com.pk/print/792938-first-ever-surgeries-to-treat-epilepsy-done-in-pakistan>

17. Wasay, M., Nadeem Saqib, M. A., & Rafique, I. (2019). Neurological Disorders Survey Pakistan 2017- 2019. Retrieved 2021, from <https://phrc.org.pk/assets/nds-2017-19.pdf>.
18. HUSSAIN A, KHAN MA, QAZI SA, RAHMAN GN. Neurology in Islamabad Children's Hospital Present Status and Problems. The Kurume medical journal. 1992 Feb 1;38(4):291-4.
19. Khatri IA, Iannaccone ST, Ilyas MS, Abdullah M, Saleem S. Epidemiology of epilepsy in Pakistan: review of the literature. Journal-Pakistan medical association. 2003 Dec 1;53(12):594-6.
20. Jawaid W, Nisa Q, Umer SR, Barry SJ, Qureshi A, Shahbaz NN. Etiology and types of seizures in patients presenting to a tertiary care hospital in Karachi: A cross-sectional study. Cureus. 2020 Jul;12(7).
21. Tahir MZ, Sobani ZA, Quadri SA, Ahmed SN, Sheerani M, Siddiqui F, Boling WW, Enam SA. Establishment of a comprehensive epilepsy center in Pakistan: initial experiences, results, and reflections. Epilepsy research and treatment. 2012;2012.
22. News T. First-ever surgeries to treat epilepsy done in Pakistan [Internet]. Today's Paper. 2021 [cited 2021Sep].
23. Sheerani M. Development of a comprehensive epilepsy surgery program in Pakistan. Journal of Pakistan Medical Association. 2005;55(1):32.
24. Khan AA, Ahmad K, Ayaz SB, Ayyub A, Akhlaq U. Cerebral palsy in Pakistani children: a hospital-based survey. Cukurova Med J. 2014 Jan 1;39(4):705-11.
25. News ARY. Lahore doctors performed successful surgery on 18 children affected by cerebral palsy [Internet]. ARY NEWS. 2021 [cited 2021Sep]. Available from: <https://arynews.tv/lahore-doctors-surgery-children-cerebral-palsy/>
26. Asif M, Younus A. Analysis of 200 Cases of Hydrocephalus Managed at DHQ Teaching Hospital Sahiwal. Pakistan Journal Of Neurological Surgery. 2018 Mar 31;22(1):21-3.
27. Bakhsh A. CSF shunt complications in infants—an experience from Pakistan. Pediatric neurosurgery. 2011;47(2):93-8.
28. Research Council PH, University Hospital AK. Neurological disorders Survey Pakistan 2017-2019 [Internet]. Neurological Disorders Survey 2017-19. 2019 [cited 2021Sep]. Available from: <https://phrc.org.pk/assets/nds-2017-19.pdf>
29. NADEEM MM, BASHIR S. Use of Endoscopic Third Ventriculostomy (ETV) Success Score to Predict the Outcome of Endoscopic Third Ventriculostomy in Obstructive Hydrocephalus. Pakistan Journal Of Neurological Surgery. 2013;17(1):35-9.
30. Ganjoo P. Challenges in pediatric neurosurgery. Journal of Neuroanaesthesiology and Critical Care. 2017 Feb;4(04): S19-23.
31. Riaz Q, Naeem E, Fadoo Z, Lohano M, Mushtaq N. Intracranial tumors in children: a 10-year review from a single tertiary healthcare center. Child's Nervous System. 2019 Dec;35(12):2347-53.
32. Khan M, Yaqoob U, Hassan Z, Uddin MM. Emergency Department Referral Profile of Traumatic Brain Injury Records at a Tertiary Care Hospital of Pakistan. Updates in Emergency Medicine. 2021 Jul 30;1(1):10-6.
33. Wani AA, Sarmast AH, Ahangar M, Malik NK, Chhibber SS, Arif SH, Ramzan AU, Dar BA, Ali Z. Pediatric head injury: a study of 403 cases in a tertiary care hospital in a developing country. Journal of pediatric neurosciences. 2017 Oct;12(4):332.

34. Bhagwati SN. Development of pediatric neurosurgery in India. *Journal of pediatric neurosciences*. 2011 Oct;6(Suppl1): S4.
35. Meara JG, Leather AJ, Hagander L, Alkire BC, Alonso N, Ameh EA, Bickler SW, Conteh L, Dare AJ, Davies J, Mérisier ED. Global Surgery 2030: evidence and solutions for achieving health, welfare, and economic development. *The Lancet*. 2015 Aug 8;386(9993):569-624.
36. Waqas M, Ujjan B, Hadi YB, Najmuddin F, Laghari AA, Khalid S, Bari ME, Bhatti UF. Cranioplasty after craniectomy in a pediatric population: single-center experience from a developing country. *Pediatric neurosurgery*. 2017;52(2):77-9.
37. Mahmood PDK. Improvement in treatment for brain tumors in Pakistan [Internet]. *The news. The News International*; 2020 [cited 2021Sep]. Available from: <https://www.thenews.com.pk/print/669526-improvement-in-treatment-for-brain-tumors-in-pakistan>
38. Abbas A, Samad L. Children at the heart of global surgery: children's surgery in low-and middle-income countries.
39. Haglund MM, Fuller AT. Global neurosurgery: innovators, strategies, and the way forward: JNSPG 75th anniversary invited review article. *Journal of neurosurgery*. 2019 Oct 1;131(4):993-9.
40. Fezeu F, Ramesh A, Melmer PD, Moosa S, Larson PS, Henderson Jr F. Challenges and solutions for functional neurosurgery in developing countries. *Cureus*. 2018 Sep;10(9).
41. Warf BC, Alkire BC, Bhai S, Hughes C, Schiff SJ, Vincent JR, Meara JG. Costs and benefits of neurosurgical intervention for infant hydrocephalus in sub-Saharan Africa. *Journal of Neurosurgery: Pediatrics*. 2011 Nov 1;8(5):509-21.